

CLAIMS

We claim:

1. An array composition comprising:
 - a) a substrate with a surface comprising discrete sites; and
 - b) a population of microspheres comprising at least a first and a second subpopulation,wherein the microspheres of each subpopulation each comprise a plurality of target analytes;
wherein said microspheres are distributed on said surface.
2. The array composition according to claim 1 wherein said microspheres of each subpopulation further comprise an optical signature.
3. The array composition according to claim 1 wherein said microspheres of each subpopulation further comprise an identifier binding ligand.
4. The array composition according to claim 3 wherein said identifier binding ligand is a nucleic acid.
5. The array composition according to claim 1 wherein said target analytes are nucleic acids.
6. The array composition according to claim 5 wherein said nucleic acids comprise genomic DNA.
7. The array composition according to claim 1 wherein said target analytes are proteins.
8. The array composition according to claim 1 wherein said substrate is a fiber optic substrate.
9. The array composition according to claim 1 wherein said substrate is plastic.
10. The array composition according to claim 1 wherein said discrete sites are wells.
11. The array composition according to claim 1, wherein said microspheres are randomly distributed on said surface.
12. The array composition according to claim 1, wherein the microspheres of said first and second subpopulation each comprise a plurality of target analytes from a first and second target source, respectively.

13. The array composition according to claim 13, wherein said first and second target source are first and second patients, respectively.

14 A method comprising:

a) providing an array composition comprising:

- i) a substrate with a surface comprising discrete sites; and
- ii) a population of microspheres comprising at least a first and a second subpopulation, wherein the microspheres of each subpopulation each comprise a plurality of target analytes;

wherein said microspheres are distributed on said surface;

b) contacting said array composition with a first set of readout probes;

c) detecting the presence of a first target analyte.

15. The method according to claim 14 further comprising:

d) contacting said array composition with a second set of readout probes;

e) detecting the presence of a second target analyte.

16. The method according to claim 14, wherein said microspheres are randomly distributed on said surface.

17. The method according to claim 14, wherein said first set of readout probes comprises at least first and second readout probes, wherein said first and second readout probes comprise first and second labels, respectively.

18. The method according to claim 17, further comprising detecting said first label as an indication of the presence of said first target analyte.

19. The method according to claim 14, wherein the microspheres of said first and second subpopulation each comprise a plurality of target analytes from a first and second target source, respectively.

20. The array composition according to claim 19, wherein said first and second target source are first and second patients, respectively.

21. A method of genotyping comprising:

a) providing an array composition comprising:

5
Sub B1

10
15
20
25

- 5
- Sub B1
- i) a substrate with a surface comprising discrete sites; and
 - ii) a population of microspheres comprising at least a first and a second subpopulation, wherein the microspheres of each subpopulation each comprise at least first and second target analytes attached to said microspheres with first and second attachment moieties, respectively;

wherein said microspheres are randomly distributed on said surface;

- 10
- b) contacting said array composition with a first set of extension probes that hybridize with at least said first target sequence adjacent to a first detection position to form an extension complex;

- c) contacting said extension complex with a composition comprising

- i) at least a first nucleotide;

- ii) polymerase;

wherein said polymerase extends a first extension probe with said first nucleotide when said first nucleotide is complementary to said first detection position of said first target sequence;

and

- c) detecting the presence of said first nucleotide.

15

22 The method according to claim 21, wherein said first nucleotide comprises a label.

20

23. A method of determining the identification of a nucleotide at a detection position in at least a first target sequence comprising:

- a) providing an array composition comprising:

- i) a substrate with a surface comprising discrete sites; and

- ii) a population of microspheres comprising at least a first and a second subpopulation, wherein the microspheres of each subpopulation each comprise a plurality of target sequences;

- 25
- b) forming a first hybridization complex between said first target sequence and at least a first readout probe; and

- c) determining the nucleotide at said detection position.

30

24. A method according to claim 23, wherein said target sequence comprises a first and a second target domain, wherein said first hybridization complex comprises said first target sequence, a first readout probe hybridized to said first domain and a second readout probe hybridized to said second domain, wherein at least one of said readout probes comprise a label said determining comprises adding a ligase to form a ligation complex.

25. The method according to claim 24, wherein said first readout probe comprises a detectable label.

26. The method according to claim 23, further comprising contacting said hybridization complex with at least a first nucleotide and a polymerase, wherein said polymerase extends said first readout probe with said first nucleotide when said first nucleotide is complementary to said first detection position of said first target sequence.

Add B2

Sub B1